

Midterm Examination
Administered: Friday, October 18, 2002

Problem (1)

Consider the first order linear ordinary differential equation.

$$\frac{dy}{dt} = f(y,t) = \sin(t)y + 2t \quad (1)$$

with the initial condition

$$y(t_0 = 0) = 1 \quad (2)$$

The second order numerical method to solve this problem is given by

$$y_i = y_{i-1} + (t_i - t_{i-1}) \frac{1}{2} [f(y_{i-1}, t_{i-1}) + f(y_i, t_i)] \quad (3)$$

- (a) Use Heun's method to approximate y at $t = 0.1$. (Use one interval of size $\Delta t = 0.1$)
- (b) Solve part (a) again but take advantage of the linearity of the ODE to avoid the approximation inherent in Heun's method.
- (c) Explain why the answers in (a) and (b) are different? Which answer is more accurate?

Problem (2)

Consider the system of linear algebraic equations:

$$\begin{aligned} x_1 + x_2 &= 2 \\ 5x_1 - 6x_2 &= -1 \end{aligned}$$

- (a) Demonstrate that the multivariate Newton-Raphson will exactly solve a system of linear algebraic equations in one iteration. Use the initial guess of your choice.

Problem (3)

Consider the system of two linear ODES.

$$\begin{aligned} \frac{dx_1}{dt} &= x_1 + x_2 \\ \frac{dx_2}{dt} &= 5x_1 - 6x_2 \end{aligned}$$

Determine the type of critical point and the stability.